

CHEKALOV, K.I.; ILLYUVIYEVA, V.P.

Use of the isotope C^{14} to study the decomposition processes of organic matter in soil. Pochvovedenie no.5:40-50 My '62.

1. Severo-Zapadnyy nauchno-issledovatel'skiy Institut sel'skogo khozyaystva. (MIRA 15:6)

(Humus) (Carbon--Isotopes)

"APPROVED FOR RELEASE: 04/03/2001

CIA-RDP86-00513R000618520008-1

APPROVED FOR RELEASE: 04/03/2001

CIA-RDP86-00513R000618520008-1"

"APPROVED FOR RELEASE: 04/03/2001

CIA-RDP86-00513R000618520008-1

APPROVED FOR RELEASE: 04/03/2001

CIA-RDP86-00513R000618520008-1"

U. S. S. R. (USSR), 1974

1974-1975 Annual Report of the

"APPROVED FOR RELEASE: 04/03/2001

CIA-RDP86-00513R000618520008-1

APPROVED FOR RELEASE: 04/03/2001

CIA-RDP86-00513R000618520008-1"

L 1741-66 ENT(1)/ENT(m)/ENP(t)/ENP(b) IJP(c) JD

ACCESSION NR: AT5013695

UR/2613/64/000/030/0093/0095

AUTHOR: Il'mas, E. R. 44.55

TITLE: The phenomenon of photon multiplication in ZnS-Mn phosphors

SOURCE AN EstSSR. Institut fiziki i astronomii. Trudy, no. 30, 1964. Issledovaniya po lyuminesentsii (Research on luminescence), 93-95

TOPIC TAGS: zinc sulfide²¹ optic material, photon multiplication, quantum yield, fluorescence yield, reflection spectrum

ABSTRACT: This is a continuation of earlier investigations by the author (with G. G. Llyd'ya and Ch. B. Lushchik, Opt. i. spektr. v. 18, 1965 and Trudy IFA AN ESSR, No. 26, 213, 1964) devoted to photon multiplication in the optical band, occurring in numerous ionic crystals such as activated alkali-halide crystals. The present study is devoted to photon multiplication in crystals of other types, particularly ionic-homopolar crystals of the ZnS type. The investigations were carried out on powdered ZnS-Mn phosphor containing 10^{-2} g/c of mangan-

Card 1/3

L 1741-66
ACCESSION NR: AT5013695

ese. The quantum-yield spectrum of this phosphor was measured with a vacuum monochromator (in the 16.7 -- 4.5 eV region) and a spectrophotometer (in the 5.9 -- 3.5 eV) region, using a procedure described in detail earlier. For comparison, the reflection spectrum of ZnS was measured with a monochromator and was found to coincide with that obtained by others, except that the structure of the individual bands was more clearly pronounced. The luminescence quantum yield spectrum showed the presence of three regions with different quantum yields. The first (3.3 -- 5.5 eV) covers the intrinsic absorption edge of ZnS and the absorption band with maximum at 3.8 eV. In the second region (5.5 -- 8.5 eV), the quantum yield varies relatively little. In the third region, beginning with 8.5 eV, the quantum yield increases sharply. An interpretation of this phenomenon was presented by the author elsewhere (with Liyd'ya and Lushchik, Opt. i spektr. v. 18, 1965). The results show that in type ZnS phosphors there occurs photon multiplication, the features of which can be explained readily by taking into consideration the features of the band structure of the crystals of this type. The author thanks Ch. B. Lushchik for direction of this work. Orig. art. has: 1 figure.

Card 2/3

L 1741-66

ACCESSION NR: AT5013695

ASSOCIATION: None

SUBMITTED: 31Oct64

ENCL: 00

SUB CODE: 01

NR REF SOV: 004

OTHER: 001

Card 3/3

The region of activator absorption, the longest wavelength of the activator

"APPROVED FOR RELEASE: 04/03/2001

CIA-RDP86-00513R000618520008-1

1. The first part of the document is a list of the names of the persons who were present at the meeting. The names are listed in alphabetical order. The names are: [illegible]

2. The second part of the document is a list of the topics that were discussed at the meeting. The topics are listed in alphabetical order. The topics are: [illegible]

3. The third part of the document is a list of the actions that were taken at the meeting. The actions are listed in alphabetical order. The actions are: [illegible]

4. The fourth part of the document is a list of the decisions that were made at the meeting. The decisions are listed in alphabetical order. The decisions are: [illegible]

5. The fifth part of the document is a list of the recommendations that were made at the meeting. The recommendations are listed in alphabetical order. The recommendations are: [illegible]

APPROVED FOR RELEASE: 04/03/2001

CIA-RDP86-00513R000618520008-1"

"APPROVED FOR RELEASE: 04/03/2001

CIA-RDP86-00513R000618520008-1

APPROVED FOR RELEASE: 04/03/2001

CIA-RDP86-00513R000618520008-1"

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

60

61

62

63

64

65

66

67

68

69

70

71

72

73

74

75

76

77

78

79

80

81

82

83

84

85

86

87

88

89

90

91

92

93

94

95

96

97

98

99

100

101

102

103

104

105

106

107

108

109

110

111

112

113

114

115

116

117

118

119

120

121

122

123

124

125

126

127

128

129

130

131

132

133

134

135

136

137

138

139

140

141

142

143

144

145

146

147

148

149

150

151

152

153

154

155

156

157

158

159

160

161

162

163

164

165

166

167

168

169

170

171

172

173

174

175

176

177

178

179

180

181

182

183

184

185

186

187

188

189

190

191

192

193

194

195

196

197

198

199

200

201

202

203

204

205

206

207

208

209

210

211

212

213

214

215

216

217

218

219

220

221

222

223

224

225

226

227

228

229

230

231

232

233

234

235

236

237

238

239

240

241

242

243

244

245

246

247

248

249

250

251

252

253

254

255

256

257

258

259

260

261

262

263

264

265

266

267

268

269

270

271

272

273

274

275

276

277

278

279

280

281

282

283

284

285

286

287

288

289

290

291

292

293

294

295

296

297

298

299

300

301

302

303

304

305

306

307

308

309

310

311

312

313

314

315

316

317

318

319

320

321

322

323

324

325

326

327

328

329

330

331

332

333

334

335

336

337

338

339

340

341

342

343

344

345

346

347

348

349

350

351

352

353

354

355

356

357

358

359

360

361

362

363

364

365

366

367

368

369

370

371

372

373

374

375

376

377

378

379

380

381

382

383

384

385

386

387

388

389

390

391

392

393

394

395

396

397

398

399

400

401

402

403

404

405

406

407

408

409

410

411

412

413

414

415

416

417

418

419

420

421

422

423

424

425

426

427

428

429

430

431

432

433

434

435

436

437

438

439

440

441

442

443

444

445

446

447

448

449

450

451

452

453

454

455

456

457

458

459

460

461

462

463

464

465

466

467

468

469

470

471

472

473

474

475

476

477

478

479

480

481

482

483

484

485

486

487

488

489

490

491

492

493

494

495

496

497

498

499

500

501

502

503

504

505

506

507

508

509

510

511

512

513

514

515

516

517

518

519

520

521

522

523

524

525

526

527

528

529

530

531

532

533

534

535

28334-66 =GT(I)/=GT(M), I/=GP(U)/ETI LJP(c) GG/AT/JP

ACC NR: AP6013073

SOURCE CODE: UR/0044/60/000/004/0064/0610

AUTHOR: Il'mas, E. R.; Lushchik, Ch. B.

ORG: Institute of Physics and Astronomy, Academy of Sciences, EstSSR (Institut Fiziki i astronomii Akademii nauk EstSSR)

TITLE: Multiplication of electronic excitations in ionic crystals /Report, Fourteenth Conference on Luminescence held in Riga 16-23 September 1968/

SOURCE: AN BSSR. Izvestiya. Seriya fizicheskaya, v. 30, no. 4, 1968, 654-660

TOPIC TAGS: crystal phosphor, semiconductor crystal, current carrier, ionic crystal, pair production, excited electron state

ABSTRACT: Each high-energy photon incident on a solid gives rise to hundreds or thousands of electron-hole pairs and other electronic excitations, which may then be manifested in different forms. The purpose of the present work was to consider the elementary processes of multiplication of electronic excitations (MEI) initiated by a UV photon that produces 2-3 excitations. Primary attention is given to the multiplication mechanism in ionic crystals. The discovery of the effect of production of two carrier pairs by one photon and other early and recent experimental studies are reviewed with numerous references to Soviet and foreign authors. The results of different investigators and the data obtained by the writers are drawn upon for more or less general discussions of photon multiplication resulting in luminescence effi-

Card 1/2

L 28334-66

ACC NR: AP6013073

ciencies exceeding unity (100%), nonlinear effects in pair production, MEE and radiation defects, MEE and photoelectronic emission, MEE in crystals of different classes, the micromechanism of electron-hole pair multiplication, and MEE involving excitons and local excitations. Illustrative energy band diagrams are adduced. In addition to effects in ionic crystals, some phenomena in semiconductors (mainly silicon² and germanium²) are described. In concluding a suggestion is made for further research. Orig. art. has: 4 figures.

SUB CODE: 20/

SUBM DATE: 00/

ORIG REF: 023/

OTH REF: 020

Card 2/2 CC

ACC NR: ATY001700

SOURCE CODE: UR/3119/66/000/004/0071/0033

AUTHOR: Il'mas, E. R.; Liyd'ya, G. G.; Lushchik, Ch. B.; Soovik, T. A.

ORG: Institute of Physics and Astronomy, AN EstSSR (Institut fiziki i astronomii AN EstSSR)

TITLE: Photon multiplication in crystals and the phenomenon of radioluminescence

SOURCE: AN LatSSR, Institut fiziki. Radiatsionnaya fizika, no. 4, 1966. Ionnyye kristally (Ionic crystals), 71-83

TOPIC TAGS: photon, radioluminescence, x ray effect, quantum yield, ionic crystal, absorption band, light excitation

ABSTRACT: In connection with their earlier experiments (Opt. i spektr. v. 18, 631, 1965 and elsewhere) dealing with observation and investigation of photon multiplication by crystals in the optical band (rather than x-ray or gamma region), the authors discuss in the present article the connection between this effect and the phenomena of x-ray luminescence and radioluminescence. Particular attention is paid to the role of different electronic excitations of the crystal lattice and to luminescence excited in ionic crystals by hard radiation. Photon multiplication in the optical range was investigated with a special set-up including a vacuum monochromator and a diffraction grating, a high power discharge lamp, a monochromator, a vacuum chamber

Card 1/2

ACC NR: AT7001786

for the samples, and a comparison standard (sodium salicylate) described in the earlier investigation. A number of optical phenomena were investigated in the photon energy range from 5 to 21 eV, particularly the spectra of the quantum yield of stationary photoluminescence of several dozen activated ion crystals. The results show convincingly that photon multiplication in the optical region of the spectrum does exist arising when a single photon produces two electronic excitations in the crystal lattice. The two possible mechanisms for this phenomenon (exciton and electron-hole) are described there and characteristic features are compared with earlier experiments by the authors and by others. It is shown that these two mechanisms operate also in the case of radioluminescence of ionic crystals. A formula is derived for the energy yield of activator luminescence excited in the main absorption bands of a crystal. The possibility of decreasing the time lag of the electron-hole radioluminescence mechanism in scintillating crystals is discussed. As a rule, in stationary radioluminescence the electron-hole mechanism predominates, while in scintillations the two mechanisms are in general on par. In NaI-Tl crystals the electron-hole mechanism apparently predominates. It is shown that a possible reason for the deviation of the real scintillation yield from the estimates presented in the article is the inertia of the electron-hole mechanism. Orig. art. has: 4 figures, 4 formulas, and 1 table.

SUB CODE: 20/ SUBM DATE: 00/ ORIG REF: 022/ OTH REF: 006

Card 2/2

IL'YAKOV, A. I.

"Tests of the Treatment of Tertian and Tropical Malaria with Biguanil", Med.
Paraz. i Paraz. Bolez., Vol. 17, No. 4, pp 312-15, 1948.

L 25603-66

ENT(m)/BWP(j) RM

ACC NR: AP6016705

SOURCE CODE: UM/0073/45/035/012/2221/2225

AUTHOR: Umayev, M. G. (Deceased) Shakirova, A. M. 24

ORG: Scientific Research Institute of Synthetic Fats (Nauchno-Issledovatel'skiy institut sinteticheskikh zhirov)

TITLE: Reaction of triphenylphosphite with omega-bromoacetophenone

SOURCE: Zhurnal obshchey khimii, v. 35, no. 12, 1965, 2221-2225

TOPIC TAGS: organic phosphorus compound, phosphinic acid, brominated organic compound, phosphate


ABSTRACT: The authors previously showed that at 160-220° trialkylphosphites, $(RO)_3P$, where $R = C_1-C_8$, undergo the Arbuzov rearrangement with omega-bromoacetophenone to form the corresponding alkyl halides and esters of ketophosphinic acids. The purpose of this article was to describe the study of the interaction of triphenylphosphite with omega-bromoacetophenone. It is shown that triphenylphosphite reacts with omega-bromoacetophenone in anhydrous benzene, toluene, or xylene or without a solvent at 120-130° to form an addition product which is thermally unstable and very reactive. It reacts readily with compounds having an active hydrogen, with the evolution of heat.

Card 1/2

UTC: 661.718.1

L 25603-66

ACC NR: AF6016705



At higher temperatures (120-140° and higher), triphenylphosphate, phenylacetylene, and hydrogen bromide are formed, which are the products of the thermal decomposition of the adduct. A mechanism for the latter reaction is proposed. [JPRS]

SUB CODE: 07 / SUBM DATE: 30Nov64 / ORIG REF: 006 / OTH REF: 007

Card 2/2 F/

1. IL'MENEV, I. S.
2. USSR 600
4. Roots (Botany)
7. Decomposition of old roots of clover, alfalfa and sainfoin, Sov. agron, 11, No. 2 1953.

9. Monthly List of Russian Accessions, Library of Congress, April 1953, Uncl.

IL'MENEV, S. I.

"Preceding Cultivation of Loamy Podzolic Soils." Sub 21 Jun 51,
All-Union Sci Res Inst of Fertilizers, Agricultural Engineering and
Soil Science.

Dissertations presented for science and engineering degrees in
moscow during 1951.

SO: Sum. No. 480, 9 May 55

IL'INEN/EV, S. I.

"The Treatment of Turf Podsollic Soils Under Grass Field Rotation Conditions." Dr Agr Sci, Moscow Order of Lenin Agricultural Acad imeni K.A. Timiryazev, Moscow, 1954. (KL, No 9, Feb 55)

SO: Sum. No. 631, 26 Aug 55- Survey of Scientific and Technical Dissertations Defended at USSR Higher Educational Institutions (14)

IL'MENEV, S. I.

Il'menev, S. I. - "The Working of Sod-Podzolic Soil under Conditions of Grass-Field Crop Rotation." Moscow Order of Lenin Agricultural Academy imeni K. A. Timiryazev. Moscow, 1956 (Dissertation for the Degree of Doctor in Agricultural Sciences).

So: Knizhnaya Letopis', No. 10, 1956, pp 116-127

IL'MENEV, S.I., prof., doktor sel'khoz.nauk

Role of agricultural practices and silvicultural measures in
controlling wind erosion. Zemledelie 23 no.8:38-45 Ag '61.
(MIRA 14:10)

1. Azovo-Chernomorskiy sel'skokhozyaystvennyy institut.
(Wind erosion)

IL'MENNY, Ye. S.

Rare metal stockworks in nepheline syenites. Izv.vys.ucheb.zav.;
geol. i razv. 1 no.11:51-58 M '58. (MIRA 12:11)

1. Moskovskiy geologorazvedochnyy institut im. S. Ordzhonikidze.
(Metals, Rare and minor)

IL'MENEV, Ya.S.; KUZIN, V.N.; NIKOL'SKIY, A.L.

Studying metamict minerals under an electron microscope. Izv. vys.
ucheb. zav.; geol. i raev. 7 no.11:126-130 N '64.

(MIRA 18:5)

1. Moskovskiy geologorazvedochnyy institut im. S. Ordzhonikidze.

IL'MENKOV, N

IL'MENKOV, N. and others. Aluminiyevaya promyshlennost': puti razvitiya v SSSR,
sostoyanie v kapitalisticheskikh stranakh. Moskva, Soi-
sekizdat, 1932. 179 p.

NN

DLC: HD9539.A614

SO: LC, Soviet Geography, Part I, 1961, Uncl.

STARIK, I.Ye.; SHEYDINA, L.D.; II' MENKOVA, L.I.

State of microquantities of radioelements in dilute solutions
Part 7: Investigation of the state of protactinium in aqueous
solutions by means of adsorption and desorption; Radiokhimiya 1
no.2:168-170 '59. (MIRA 12:8)
(Protactinium) (Sorption)

STARIK, I.Ye.; SHEYDINA, L.D.; IL'MENKOVA, L.I.

State of microquantities of radioelements in solutions. Part 10:
Study of the state of protactinium in aqueous solutions. Radiokhimiia
1 no.4:391-394 '59. (MIRA 13:1)
(Protactinium)

J. L. M. N. K. O. U. T. L. I.

WINZBURG, F.L.

21 (a). 5 (a)

ATTENTION

17-00000

Practical:

STYLING

2017/09-7-3-17/24

Michaletsky, Y. I.

All-Union Symposium on Radiochemistry (Vsesoyuzny simpozium po radiokhimii)

Atomnaya energiya, 1959, Vol 7, Nr 2, pp 179-176 (ISSR)

[illegible]

Quest 1/3

☆

IL'MENKOVA, L. I., Cand Chem Sci -- (diss) "State of protactinium in aqueous solutions." Leningrad, 1960. 17 pp with graphs; (Academy of Sciences USSR, Radium Inst im V. G. Khlopin); 250 copies; price not given; (KL, 22-60, 132)

S/186/61/004/001/006/020
A051/A129

21,3100

AUTHORS: Sheydina, L.D., Il'menkova, L.I.

TITLE: A study on the state of protactinium in aqueous solutions by the extraction method

PERIODICAL: Radiokhimiya, v 3, no 1, 1961, 24-30

TEXT: The authors have used the extraction method for studying the state of protactinium in aqueous solutions. Tri-n-butylphosphate (TBPh) in benzene was used as the extracting agent. The artificial Pa^{233} isotope was used for the experiments, and it was shown that at a pH of the solution equal to 0-2 the predominating state is the ion of average composition, $\text{Pa}(\text{OH})_{4.5}^{0.5+}$. With an increase in the pH of the solution the formation of more complex ions with a low positive charge takes place. At pH=5 the formation of protactinium hydroxide $\text{Pa}(\text{OH})_5$ begins. The data obtained in the investigation of the Pa^{233} extraction with a 25% mixture of TBPh-benzene from solutions of various pH (from 0 to 9) agree with previous assumptions of the authors (Ref 1-6).

Card 1/8

23073

S/186/61/003/001/006/020

A051/A129

A study on the state of protactinium ...

With an increase in the concentration of the nitric acid (1-14 n HNO_3) an increase in the content of neutral forms of protactinium occurs. It is assumed that a change in the composition of these neutral forms of protactinium takes place when changing from 1-3 n HNO_3 to 3-5 n concentration. In the experimental procedure the apparent distribution coefficient (α_{app}) was computed as the ratio of equilibrium concentrations of protactinium in the organic and aqueous phases, $\alpha_{\text{app}} = \frac{C_{\text{org.}}}{C_{\text{aqu.}}}$, and the % extraction to the initial concentration of protactinium in the aqueous phase multiplied by 100:

$$\% \text{ extraction} = \frac{C_{\text{org.}}}{C_{\text{initial aqueous}}} \cdot 100,$$

where $C_{\text{org.}}$ is the concentration of Pa^{233} in the organic phase, $C_{\text{initial aqueous}}$ the initial concentration of Pa^{233} in the aqueous phase. The low percentage of extraction of Pa^{233} with a drop in the concentration of HNO_3 is explained either by the shortage of NO_3^- ions needed for binding Pa into a neutral complex or by a deeper hydrolysis of the Pa compounds. In order to determine the


Card 2/8

73873

8/186/61/003/001/006/020
A051/A129

A study on the state of protactinium ...

forms of protactinium within the pH-1 range, the authors conducted the following mathematical analysis: Protactinium in the aqueous solution is considered to be in the form of nitratehydroxo-complexes of various composition, the formation of which takes place according to Formula 1:



The sum constant of formation of these complexes ($K_{\text{formation}}$) is expressed through Formula 2:

$$K_{\text{form.}} = \frac{\left[\sum_n \sum_m \text{Pa}(\text{OH})_n(\text{NO}_3)_m^{5-(n+m)} \right]}{\left[\sum_n \text{Pa}(\text{OH})_n^{5-n} \right] \cdot [\text{NO}_3^-]^m} \quad (2)$$

Since Pa is extracted by TBPh in the form of a neutral complex of the type:

Card 3/8


23073

S/186/61/003/OC1/006/020

AO51/A129

A study on the state of protactinium ...

$\text{Pa}(\text{OH})_n(\text{NO}_3)_k \cdot x\text{TBPh}$ (Ref 6,7) where $n+k=5$, the extraction equation is expressed by the reaction:



The distribution constant of reaction (3) is determined by Formula 4:

$$K_T = \frac{\left[\sum_n \sum_k \text{Pa}(\text{OH})_n(\text{NO}_3)_k \cdot x\text{TBPh} \right]_{\text{org.}}}{\left[\sum_n \sum_m \text{Pa}(\text{OH})_n(\text{NO}_3)_m^{5-(n+m)} \right] \cdot [\text{TBPh}]^x \cdot [\text{NO}_3]^{5-(n+m)}} \quad (4)$$

where $n+k=5$, or substituting from equation (2), equation (5) is derived:

Card 4/8

23473

S/186/51/004/001/006/020
A051/A129

A study on the state of protactinium ...

$$K_r = \frac{\left[\sum_n \sum_k \text{Pa}(\text{OH})_n (\text{NO}_3)_k \cdot x \text{TBPh} \right]_{\text{org.}}}{K_{\text{org.}} \cdot \left[\sum_n \text{Pa}(\text{OH})_n^{5-n} \right] [\text{NO}_3^-]^{5-n} [\text{TBPh}]^x} \quad (5)$$

On the other hand the distribution coefficient ($\alpha_{\text{app.}}$) is considered to be the ratio of the Pa concentration in the organic phase to the Pa concentration in the aqueous phase (equation 6):

$$\alpha_{\text{app.}} = \frac{C_{\text{org.}}}{C_{\text{aqu.}}} \quad (6)$$

Equation 16: $\lg \alpha_{\text{app.}} = (5-n) \lg [\text{H}^+] + \lg \text{const.} \quad (16)$

is used to determine (5-n) from the slope of the curve on the relationship graph: $\lg \alpha_{\text{app.}}$ versus $\lg \text{H}^+$, where n is the average number of OH⁻ groups.

Card 5/e

1073

S/186/41/001/001/006/020
A051/A129

A study on the state of protactinium ...

bound with one atom of Pa (Fig 2). The authors point out that this mathematical analysis can only serve to judge the average composition of the Pa forms and is not a quantitative evaluation. Fig 3 shows the relationship of the % extraction of Pa with respect to the HNO_3 concentration. A study of the logarithmic relationship of the distribution coefficient and the activity of HNO_3 leads the authors to assume that within the concentrations range of 1-5 n at least two forms of neutral Pa molecules are present. There are 4 graphs, 16 formulae and 12 references: 9 Soviet-bloc, 4 non-Soviet-bloc.

Card 6/8

22995

S/186/61/003/002/005/018
E037/E419

21.3230

AUTHORS: Starik, I.Ye., Sheydina, L.D. and Il'menkova, L.I.

TITLE: The state of protoactinium in aqueous solution
IV. Ion-exchange study

PERIODICAL: Radiokhimiya, 1961, Vol.3, No.2, pp.150-154

TEXT: C.J.Hardy, D.Scargill and J.M.Fletcher (Ref.9: J.Inorg. Nucl.Chem.7, 3, 257 (1958)) have studied the adsorption of protoactinium (10^{-4} - 10^{-7} M) on De-Acidite FF and Zeocarb 225 from nitric acid solution. This work is concerned with the ion-exchange behaviour of micro-quantities of Pa^{233} in nitric acid solutions and with determination of the charges on the cations and anions in the HNO_3 concentration range studied. A batch procedure was used with 10^{-11} M Pa^{233} solutions and 200 mesh Dowex 50 (H^+ -form) and Dowex 1 (NO_3^- -form). The distribution coefficient K_p was calculated as the ratio of the Pa^{233} activity adsorbed by 1 g of resin to the equilibrium activity of 1 ml of solution, i.e.

$$K_p = \frac{(A_1 - A_2) V}{pA_2}$$

where A_1 is the specific activity of the initial solution,
Card 1/6

22995

The state of protoactinium ...

S/186/61/003/002/005/018
E037/E419

A_2 is the specific activity of the solution after equilibration with the resin, V is the volume of solution and p the weight of resin. Fig.1 and 2 show the Pa^{233} distribution coefficient as a function of nitric-acid concentration for the cation-exchange and anion-exchange resins used. It is evident that positively-charged Pa species exist in 1-5 N HNO_3 . The increase in K_p above 5 N HNO_3 , as shown in Fig.2, reflects the increasing charge on the Pa anion complexes. The slight increase in K_p above 10-12 N HNO_3 which is observed for the anion-exchange resin (Fig.1) is probably due to the conversion of hydroxy groups into equo-groups. J.D.Strickland's (Ref.15; Nature, 169, 620 (1952)) method was used to determine the charge on the protoactinium ions. The charge on an ion is given by the slope of a plot of $\log C_1/(C_2 - C_1)$ vs. $\log [\text{H}^+]$ or $\log [\text{NO}_3^-]$, where C_1 is the concentration of the ion of unknown charge in the aqueous solution after equilibration with the resin and C_2 is the initial concentration of the same ion. The slope of the $\log [\text{H}^+]$ dependence is found to be 2.6 ± 3 from Fig.3, so that the mean charge on the Pa^{233} cation in the resin phase (1-4 N HNO_3) is 3. The slope of the $\log [\text{NO}_3^-]$ dependence is 3.1 (Fig.4) and this corresponds to the mean charge on the

Card 2/6

22995

S/186/61/003/002/005/018

E037/E419

The state of protoactinium ...

protoactinium anion in the resin phase (8 - 12 N HNO_3). It is clear that in HNO_3 solutions of these concentrations Pa ions with charges +3 and -3 and below exist. Our results do not establish the numbers of OH^- and NO_3^- coordinated to Pa but the following forms are possible:

in 1 - 4 N HNO_3 : $\text{Pa}(\text{OH})_3^{3+}$; $[\text{Pa}(\text{OH})_2(\text{NO}_3)]^+$; $\text{Pa}(\text{OH})_2^{2+}$; $[\text{Pa}(\text{OH})_2(\text{NO}_3)]^{2+}$;
 $[\text{Pa}(\text{OH})_2(\text{NO}_3)_2]^+$; $\text{Pa}(\text{OH})_2^+$.

in 8 - 12 N HNO_3 : $[\text{Pa}(\text{OH})(\text{NO}_3)_2]^-$; $[\text{Pa}(\text{OH})(\text{NO}_3)_3]^{2-}$; $[\text{Pa}(\text{OH})(\text{NO}_3)_4]^{3-}$;
 $[\text{Pa}(\text{NO}_3)_3]^-$; $[\text{Pa}(\text{NO}_3)_4]^{2-}$; $[\text{Pa}(\text{NO}_3)_5]^{3-}$.

The neutral forms of protoactinium have not been considered in this article; they were described by the authors in a previous paper (Ref.14: Radiokhimiya, 3, 1, 24 (1961)). The data obtained by the authors agree with and complete those given by C.J.Hardy et al (Ref.9). There are 4 figures and 16 references: 7 Soviet-bloc and Card 3/6

22995

The state of protoactinium ...

S/186/61/003/002/005/018
E037/E419

X

9 non-Soviet-bloc. The four most recent references to English language publications read as follows: A.G.Maddock, J.Inorg.Nucl. Chem., 2, 2, 114 (1956); G.R.Choppin, J.Chem.Ed., 36, 9, 462 (1959); C.J.Hardy, D.Scargill, J.M.Fletcher, J.Inorg.Nucl.Chem., 7, 3, 257, (1958); K.A.Kraus, D.C.Michelson, F.Nelson, J.Am.Chem.Soc., 81, 13, 3204 (1959).

SUBMITTED: April 18, 1960

Card 4/6

STARIK, I.Ye.; SHEYDINA, L.D.; IL'MENKOVA, L.I.

State of protactinium in aqueous solutions. Part 6: Adsorption
properties of protactinium. Radiokhimiya 4 no.13:4-9 '62.
(MIRA 15:4)

(Protactinium) (Adsorption)

STARIK, I.Ye.; SHEYDINA, L.D.; IL'MENKOVA, L.I.

Study of the state of protactinium in aqueous solutions. Part 5:
Region of occurrence of pseudocolloids. Radiokhimiya 3 no.6:690-
693 '61. (MIRA 14:12)

(Protactinium)
(Colloids)

STANIK, I.I.; IL'MENKOVA, I.I.

State of the microquantities of elements in aqueous solutions.
Cation complexes of protactinium. Radiokhimiia 5 no. 6:
679-683 '63. (MR: 1717)

AVDEYEV, A.I., inzh.; IL'MENSKIY, D.A., inzh.; NOTARIUS, M.D., inzh.

Using asymmetrical bridge circuits in magnetoelectric ratiometers
of resistance thermometers. Priborostroenie no.6:1-3 Je '61.
(MIRA 14:6)

(Bridge circuits)
(Thermometers)

IL'MER, A.L., inzh.

Heat protection of transformers in mines. Bezop.truda v prom.
4 no.2:29 F '60. (MIRA 13:5)

1. Trest Kirgisugol'.
(Electricity in mining--Safety measures)

IL'MINSKAYA, R. I.

"The Interpollination of Basic Apple Varieties in Omskaya Oblast and the Order of Their Arrangement in the Orchard." Cand Agr Sci, Omsk Agricultural Inst imeni S. M. Kirov, Omsk, 1954. (KL, No 7, Feb 55)

SO: Sum. No. 631, 26 Aug 55-Survey of Scientific and Technical Dissertations Defended at USSR Higher Educational Institutions (14)

PETRENKO, V.G.; SEMISALOVA, V.N.; Prinimala uchastiye Il'minskaya, V.I.

Coking blended coal charges with petroleum residue additions
and coal tar. Koks i khim. no.16:14-17 '61. (HIRA 15:2)

1. Orsko-Khalilovskiy metallurgicheskiy kombinat.
(Coke industry)

S/194/61/000/006/071/077
D201/D302

AUTHOR: Il'minskiy, N.Ya. and Loyter, Ye.G.
TITLE: Junction transistor frequency divider
PERIODICAL: Referativnyy zhurnal. Avtomatika i radioelektronika,
no. 6, 1961, 12, abstract 6 K81 (V sb. Poluprovodnik.
pribory i ikh primeneniye, no. 5, M., Sov. radio,
1960, 254-263)

TEXT: The principle of operation is considered of a junction transistor HF divider (D) together with the results of experimental analysis of the steady state operation of D with junction transistor П14 (P14) with division factor 5. The operation of D was examined at a frequency of 1 Mc/s in a common emitter circuit. The divider can also work in common base configuration, the circuit has been found to be, however, less stable in operation. D operates also with other than P14 transistors, provided the current gain cut-off frequency is several times higher than the output frequency from the

Card 1/2

Junction transistor...

S/194/61/000/006/071/077
D201/D302

divider. 4 references. [Abstracter's note: Complete translation]

✓

Card 2/2

IL' MINSKIY, V. Ya.
USSR/Engineering - Instrumentation

FD-2615

Card 1/1 : Pub. 41-1/21

Author : Il'minskiy, V. Ya., Moscow

Title : Investigation of the elasticity characteristics of diaphragms

Periodical : Izv. AN SSSR, Otd. Tekh. Nauk 4, 3-21, Apr 1955

Abstract : Presents results of experimental investigation of the characteristics of diaphragms with simple sinusoidal and angular corrugations. The relationships thus obtained are sufficiently accurate for determining the characteristics of diaphragms of a given cross section. Analyzes the results of the investigation and compares theoretical and experimental relationships. Presents a short review of contemporary methods for computing corrugated diaphragms. Described experimental methodology. Formulae, graphs, tables. Eight references, 5 USSR

Institution :

Submitted : December 28, 1954

AUTHOR: Il'minskiy, V. Ya. (Moscow)

SOV/24-58-8-22/37

TITLE: On Extending the Limit of Application of Nomograms for
Calculating Diaphragms with Angular Goffering
(O rasshirenii predela ispol'zovaniya nomogram dlya
rascheta membran s uglovym gofrom)

PERIODICAL: Izvestiya Akademii Nauk SSSR, Otdeleniye Tekhnicheskikh
Nauk, 1958, Nr 8, pp 120-123 (USSR)

ABSTRACT: In earlier work of the author an equation was derived
for calculating the characteristics of diaphragms with
sinusoidal and angular goffering, which is suitable for
calculating diaphragms of 24 to 60 mm dia. To extend the
limits of applicability of the experimentally determined
relations it was decided to apply the law of analogy,
according to which the bending in geometrically similar
diaphragms should be proportional to their dimensions.
This law follows directly from a dimensional analysis of
the theoretical formulae (see Panov, Ref 2 and Fedosev
Ref 3) which can be written in the form expressed by
Eq.(1). It follows from the derived relation, Eq.(2),
that the bending of the diaphragms should change in

Card 1/4 proportion to their dimensions and the authors considered

BOV/24-58-8-22/37

On Extending the Limit of Application of Homograms for
Calculating Diaphragms with Angular Goffering

it necessary to verify this law of proportionality of the characteristics for real diaphragms. The verification was carried out on diaphragms with angular goffers, Fig.1, made of beryllium bronze BrB 2.5. The technology of manufacture and determination of its characteristics were the same as in the earlier published work of the author (Ref 1). The dimensions of the diaphragms and the measured bending values are entered in Tables 1 and 2, where N denotes the number of the investigated diaphragms. The characteristics of diaphragms with various radii were reduced to those of a diaphragm with a 24 mm radius by changing the scale of the bending values in accordance with the similarity factor entered in Fig.2. The numerical values of these data are entered in Table 3 and it can be seen that in similar diaphragms the deviations in the characteristics from the proportionality law does not exceed on the average + 7.7% of the current and $\pm 2.5\%$ of the maximum bending value of the diaphragm. The fact that there is some divergence in the characteristics is attributed to imperfections in the manufacture

Card 2/4

SOV/24-58-8-22/37

On Extending the Limit of Application of Nomograms for
Calculating Diaphragms with Angular Goffering

of the diaphragms. It follows from the derived relations, Eq.(8), p 123, that the stresses in similar diaphragms and consequently also the permissible pressures should be equal. Experimental determination of the permissible pressure in the investigated diaphragms has shown that in all cases irrespective of the diameter of the diaphragm they are within the limits of 3.8 to 4.2 kg/cm². The derived relations permit extending considerably the range of utilisation of the graph published in the previous paper which was made for determining the permissible pressure on a diaphragm of 48 mm dia. In order to determine the permissible pressure on a diaphragm of any diameter, the dimensions of that diaphragm have to be reduced to 48 mm by multiplying with the proportionality coefficient and then to determine by means of the graph the permissible pressure which will equal the sought pressure. A nomogram is included (Fig.4) for calculating diaphragms with angular goffers; by means of this diagram it is also possible to solve the inverse problem, namely, to determine the dimensions of the diaphragms on the basis of specified

Card
3/4

SOV/24-58-8-22/37

On Extending the Limit of Application of Nomograms for
Calculating Diaphragms with Angular Goffering

characteristics. A calculation example is included. Acknowledgments are made to Professor S. S. Tikhmenev for his criticism and advice about the manuscript and also to Engineer N. I. Taruntayeva and M. P. Luzhenkov for their assistance in carrying out the experiments. There are 4 figures, 3 tables and 3 Soviet references.

SUBMITTED: November 12, 1957

1. Diaphragms (Mechanics)--Deflection
2. Diaphragms (Mechanics)--Stresses
3. Diaphragms (Mechanics)--Mathematical analysis
4. Nomographs--Applications

Card 4/4

CATEGORY : Farm Animals.Honeybees
ASS. JOUR. : RZBiol., No. 13 1958, No. 59654
AUTHOR : Il'mukhin, N.D.
INST. :
TITLE : How Did I Save an Apiary From the Foul Brood?

ORIG. PUB. : Pchelovodstvo, 1957, No.11, 52-53

ABSTRACT : The use of sodium norsulfazole and sulcymide proved ineffective. The apiary was sanitized with the aid of penicillin which was fed in sugar syrup to the sick honeybees in a dose of 200,000 I.U. and to the healthy ones in a dose of 100,000 I.U.

CARD: 1/1

1. P. D. IL'YUKHIN
2. USSR (600)
4. Bee Culture - Queen Rearing
7. Obtaining fertile queens. Pchelovodstvo 29 no. 12. 1952.
9. Monthly List of Russian Accessions, Library of Congress, April 1953, Uncl.

ILMURZYNSKA, Krystyna; ZAREMBA, Janusz

A case of isolated myocarditis in pregnancy and puerperium. Pol.
tyg. lek. 17 no.5; 185-188 29 Ja '62.

1. Z Kliniki Kardiologii SDL AM w Warszawie; kierownik: prof. dr
med. Edmund Zera i z Pracowni Anatomopatologicznej; kierownik:

J. Zaremba — Szpitala Miejskiego Nr 6 Warszawa.

(MYOCARDITIS in pregn) (PREGNANCY compl)

(PUERPERIUM compl)

SZACHOWSKI, J.; ILMURZYNSKA, K.

Effect of amyl nitrite on cardiac murmurs. Kardiol. Pol. 7
no.2:103-108 '64.

1. Z Kliniki Kardiologii Studium Doskonalenia Lekarzy Akademii
Medycznej w Warszawie (Kierownik: prof. dr E. Pera).

IIMURZYNSKA, Krystyna

Late systolic sounds in normal subjects. Pol. tyg. lek. 19 no. 47:
1817-1819 23 N'64.

1. Z Kliniki Kardiologii Studium Doskonalenia Lekarzy w Akademii
Medycznej w Warszawie (kierownik: prof. dr. med. E. Zera).

GOROWSKI, Tadeusz; ILMURZYNSKA, Krystyna

Phonothyrography — a new method for the examination of thyroid blood supply. Pol. tyg. lek. 17 no 418 577-1582 8 0 '62.

1. Z I Katedry Choroby Wewnętrznych; kierownik: prof. dr med. W. Hartwig
i z Katedry Kardiologii, kierownik: prof. dr med. E. Zera — Studium
Doskonalenia Lekarzy AM w Warszawie.
(AUSCULTATION) (THYROID GLAND)

ILMURZYNSKA, K.

Selection of filters for phonocardiography. Kardiol. pol. 6
no.2:105-112 '63.

1. Z Kliniki Kardiologii Studium Doskonalenia Lekarzy AM w
Warszawie Kierownik: prof. dr E. Zera.
(PHONOCARDIOGRAPHY) (FILTERS)

ILMURZYŃSKA, Krystyna; SIDOROWICZ, Wacław

Attempted evaluation of a phonocardiographic phenomenon in athletes. Pol. tyg. lek. 18 no.13:467-470 25 Mr '63.

1. Z Kliniki Kardiologii Studium Doskonalenia Lekarzy A.M. w Warszawie; kierownik: prof. dr med. E. Żera i w Centralnej Przychodni Sportowo-Lekarskiej.
(PHONOCARDIOGRAPHY) (SPORT MEDICINE)

ILMURZYNSKA, K.; SZACHOWSKI, J.

Effect of phenylpinephrine. Kardiolog. pol. 6 no. 4: 233-239 '63.

1. Z Kliniki Kardiologii SMD w AM w Warszawie; kierownik:
prof. dr. E. Zera.

*

ZERA, Edmund; HOFFMAN, Maria; JANIĆ, Zofia; ILMURZYŃSKA, Krystyna;
KRZYŻANOWSKA, Regina

Rehabilitation of myocardial infarction patients under
sanatorial conditions. Pol. tyg. lek. 18 no, 34:1264-1267
19 Ag 1963.

1. Z Kardiologicznego Ośrodka Rehabilitacji Pospitalnej w
Sanatorium w Maleczie i z Kliniki Kardiologii Studium
Doskonalenia Lekarzy w AM w Warszawie; kierownik: prof. dr
med. Edmund Zera.

(MYOCARDIAL INFARCT) (REHABILITATION)

Ilmurzynski, E.

"A different approach to selective thinning."

p. 7 (Sylvan, Vol 102, no. 9, Sept 1958, W_arsaw, Poland)

Monthly Index of East European Accessions (AAEI) LC, Vol 8, No. 1, Jan 59.

ILMURZYNSKI, Eugeniusz

Trends in silviculture in Poland. Sylwan 104 no.4:55-61.
Ap '60.

POLAND

ILNURZYNSKA, Krystyna and SIDOROWICZ, Wacław, Cardiology Clinic (Klinika Kardiologii), Physicians' Post-graduate Courses (Studium Doskonalenia Lekarzy) of the AM [Akademia Medyczna, Medical Academy] in Warsaw (Director: Prof. Dr. med. B. ZERA) and Central Sports-Medical Outpatient Clinic (Centralna Przychodnia Sportowo-Lekarska)

"Phonocardiographic Tracings in Sportsmen."

Warsaw, Polski Tygodnik Lekarski, Vol 18, No 13, 25 Mar 63, pp 467-470.

Abstract: [Authors' English summary] Phonocardiographic tracings were performed in 30 bicyclists of the national olympic cadre. No abnormal tracings were noted. The value of phonocardiographic tracings in the medicine of sportsmen, especially in the interpretation of systolic murmurs, is discussed. There are seven (7) references, of which one (1) each are Polish, Russian, and Italian, and the remaining four (4) are English.

1/1

IL'NITSKAYA, G.D. [il'nyts'ka, H.D.]


Prospects for the use of substitutes of food raw materials for
technical purposes. Khar. prom. no.3:61-62 J1-S '65. (MTRA 18:9)

IL'NITSKIY, Iosif Ivanovich; KHODAKOVSKIY, N.S., inzh., red.;
BOGOSLAVETS, N.P., tekhn. red.

[Automatic and semiautomatic machine tools] Stanki-avtomaty i
poluavtomaty. Moskva, Gos. nauchno-tekhn.izd-vo mashinostroit.
lit-ry, 1961. 46 p. (Nauchno-populiarnaya biblioteka rabochego
stanochnika, no.30) (MIRA 15:3)
(Machine tools) (Automatic control)

A capacitive feedback differentiating... ⁵³¹⁷⁸ S/103/b2/023/001/009/014
D201/D304

Fig. 1.



Card 3/3

CHITSKAYA, N. M.

New species of Middle Sarmatian Ostracoda in the southern Ukraine.
Paleont. zhur. no. 3: 135-139 '64. (MIRA 18:2)

1. Odesskiy gosudarstvennyy universitet.

IL'NITSKAYA, N.M.

Ostracoda from the clayey facies of Pontic deposits (underlying the limestone) of the Black Sea shore near Odessa. Paleont. zhurn. no.2: 107-108 '59. (MIRA 13:1)

1. Odesskiy gosudarstvennyy universitet, Paleontologicheskiy muzey.
(Odessa region--Ostracoda, Fossil)

IL'NITSKAYA, N.M.

Middle Sarmatian ostracods from the environs of Odessa. Paleont.
zhur. no.4:135-136 '60. (MIRA 14:1)

1. Odesskiy gosudarstvennyy universitet, Paleontologicheskii muzey.
(Odessa region--Ostracoda, Fossil)

USSR / Microbiology. Microbes Pathogenic to Man
and Animals. Corynebacteria.

F

Abs Jour : Ref. Zhur - Biol., No. 21, 1958, No 95195

Author : Il'nitskaya, Ya. A.

Inst : Odessa Scientific-Research Institute of
Epidemiology and Microbiology.

Title : Materials for the Problem of Obtaining High
Quality Diphtheria Anatoxins. Report I.
Study of the Possibility of Isolating Highly-
Toxic Cultures of a Diphtheria Bacillus From
a Productive Strain of the PW8 Diphtheria
Microbe.

Orig Pub : Tr. Odessk. n.-i. in-ta epidemiol. i mikrobiol.
1957, 3, 103-109

Abstract : The capacity was determined for toxic form-
ation of 298 subcultures of the R-form of

Card 1/2

USSR / Microbiology. Microbes Pathogenic to Man
and Animals. Corynebacteria.

F

Abs Jour : Ref. Zhur - Biol., No. 21, 1958, No. 95195

the PW₈ strain in Martin's broth which contained 0.25% glucose. Toxicity was varied in the subcultures studied, and of 10 subcultures selected on the basis of increased toxic formation, only in 2 was this indication stable. Biostimulators of growth (aloe, FIBS and peloidin) which were added to the nutrient medium in the quantity of 0.1 and 0.5% decreased the toxicity of the cultures or had no effect on them.

Card 2/2

IL'NITSKAYA, Ye.A., Cand Med Sci -- (diss) "Diphtheria antitoxins and
stability of their antigenic and immunogenic properties in
storage." Odessa, 1959, 11 pp (Odessa State Med Inst im N.I. Pirogov)
200 copies (KL, 36-59, 119)

- 95 -

IL'NITSKAYA, Ye.I.

Determination of coal resistance to shifting and faulting. Truly
Inst.gor.dela 1:124-135 '54 (MLRA 7:12)
(Coal geology) (Faults (Geology))

IL'NITSKAYA, YE. I.

IL'NITSKAYA, YE. I. -- "Methods of Determining the Resistance of Coal to Shearing and Fracture." Acad Sci USSR, Inst of Mining, Moscow, 1955. (Dissertations for the Degree of Candidate of Technical Sciences)

SO: Knizhnaya Letopis' No. 39, 24 Sept 55

"APPROVED FOR RELEASE: 04/03/2001

CIA-RDP86-00513R000618520008-1

APPROVED FOR RELEASE: 04/03/2001

CIA-RDP86-00513R000618520008-1"

IL'NITSKAYA, Ye. I.

"Mechanical Extraction of Coal."

report presented at the Conference in the Mining Inst. AS USSR on Problems
of Rock Disintegration, 20-22 May 1958.
(Vest. AN SSSR, No. 8, 1958, pp. 130-132)

PROTOD'YAKONOV, M.M., doktor tekhn.nauk, prof.; IL'NITSKAYA, Ye.I., kand.
tekhn. nauk

New techniques in coal mining. Bezop. truda v prom. 2 no. 6:19-
21 Jo '58. (MIRA 11:7)

(Coal mines and mining)

AUTHORS: Il'nitskaya, Ye.I., Candidate of Technical Sciences SOV-127-58-8-23/27

TITLE: All-Union Conference on Problems of Rock Crushing (Vsesoyuznoye soveshchaniye po problemam razrusheniya gornyx porod)

PERIODICAL: Gornyy zhurnal, 1958, Nr 8, p 74 (USSR)

ABSTRACT: The above mentioned conference was convened on 20 - 23 May 1958 and took place in the Institute of the Mining Industry AS USSR, in Moscow. Representatives of 82 scientific institutes of the AS USSR and of Academies of Sciences of other republics, as well as representatives of vases, projects institutions, ministries and industrial organizations took part. The senior scientific collaborator of the VNIIZhelezobeton, N.K. Timchenko, demonstrated a mechanical indicator of the strength of stone. A.A. Pavlikov (Tomskiy politekhnicheskii institut) (the Tomsk Polytechnical Institute) and A.I. Gol' - binder (Vsesoyuznyy nauchno-issledovatel'skiy institut burovoy tekhniki) (the All-Union Scientific Research Institute of Drilling Technique) reported on new methods of jet and explosion drilling. Generalized rules of percussive, rotating and rotating-percussive drilling were reported on by Professor M.I. Koyfman, (IGD AS USSR). M.I. Ryngel's (VNIINT), Candi-

Card 1/2

All-Union Conference on Problems of Rock Crushing

SOV-127-58-8-23/27

date of Technical Sciences reported on new methods of rock crushing. Engineer N.G. Kartavy (The Moscow Mining Institute) demonstrated the method of rock and coal crushing described by M.B. Gordon. R.V. Akopov (Institut stroymaterialov i sooruzheniy Armyanskoy SSR) (The Building Material and Constructions Institute of the Armenian SSR) reported on the wear and tear on drilling cutters. Professor M.M. Protod'yakonov and Engineer B.M. Loguntsov (IGD AS USSR) reported on the generalization of more than 60 scales of resistance to drilling and on the comparison of these scales with the coefficients of rock strength. The Candidate of Technical Sciences I.G. Malikidze (IGD of AS of Georgian SSR) presented an improved method for determining the strength of rocks (first developed by Protod'yakonov and V.S. Voblikov). The conference decided to continue research on all discussed problems to develop a general system of mechanical rock crushing.

ASSOCIATION: (IGD AS USSR) (Mining Institute, AS USSR)

1. Rock--Processing
2. Coal--Processing
3. Mines--Equipment
4. Mining engineering

Card 2/2

(5)(4)

IL'NITSKAYA, Ye.I.

Effect of size on the strength of rocks. Fiz.-mekh.svois., dav.i
razr.gor.porod no.1:17-23 '62. (MIRA 16:3)
(Rocks--Testing)

KOYFMAN, Mikhail Il'ich; IL'NITSKAYA, Yelena Ivanovna; KARPOV,
Viktor Ivanovich; PROTOD'YAKONOV, M.M., prof., doktor
tekhn. nauk, otv. red.; TEDER, R.I., otv. red.

[Resistance of rocks in a volume stressed state; some
problems in the methodology of research] Prochnost' gor-
nykh porod v ob'emnom napriazhennom sostoianii; nekotorye
voprosy metodiki issledovani. Moskva, Nauka, 1964. 32 p.
(MIRA 17:11)

BRESLAV, I.S.; ZHIRONKIN, A.G.; IL'NITSKIY, A.M.; KONZA, E.A.;
MITIUSHOV, M.I.; NOZDRACHEV, A.D.; SALATSINSKAYA, Ye.N.;
TROSHIKHIN, G.V.; SEMELEVA, A.M.

Some data on the effect of a closed space on the physiological
functions in animals. Probl.kosm.biol. 2:291-302 '62.

(MIRA 16:4)

(SPACE MEDICINE)

SECRET

CONFIDENTIAL

CONFIDENTIAL

CONFIDENTIAL

CONFIDENTIAL

CONFIDENTIAL

CONFIDENTIAL

"APPROVED FOR RELEASE: 04/03/2001

CIA-RDP86-00513R000618520008-1

SECRET
TOP SECRET

APPROVED FOR RELEASE: 04/03/2001

CIA-RDP86-00513R000618520008-1"

ACC NR: AP6021592

(N)

SOURCE CODE: UR/0402/66/000/003/0374/0374

AUTHOR: Il'nitskiy, A. P.

ORG: Department of Community Hygiene, First Moscow Medical Institute (Kafedra kommunal'noy gigiyeny Pervogo Moskovskogomeditsinskogo instituta im. I. M. Sechenova); Laboratory for Etiology and Diagnostics of Acute Respiratory Infections, Institute of Virology, Academy of Medical Science, SSSR, (Laboratoriya etiologii i diagnostiki ostrykh respiratornykh infektsiy Instituta virusologii im. D. I. Ivanovskogo, AMN SSSR)

TITLE: Water and the dissemination of adenoviruses

SOURCE: Voprosy virusologii, no. 3, 1966, 374

TOPIC TAGS: epidemiology, adenovirus, human disease, aerosol, *WATER POLLUTION, VIRUS*

ABSTRACT:

It is possible that adenoviruses can contaminate cisterns and waterways besides infecting by more usual routes. Adenovirus types 3, 4, and 7a survived more than 20 days in the laboratory at 19-22°C and pH 7.4.

Users of contaminated water were infected with keratoconjunctivitis although adenoviruses have not yet been isolated from water sources.

[W.A. 50; CBE No. 10]

SUB CODE: 06/ SUBM DATE: none/

Card 1/1

IL'NITSKIY, D., inzh. (Kiyevskaya oblast')

Device for removing roller bearings. Sel'.mekh. no.3:32-33 '62.
(MIRA 15:3)

(Tractors--Equipment and supplies)

DOVGICH, I.O. [Dovhich, I.O.], starshiy nauchnyy sotrudnik; IL'NITSKIY, D.V.
[Il'nyts'kyi, D.V.], mladshiy nauchnyy sotrudnik

Improve the care of the lubrication system. Mekh. sil'. hosp. 13
no.9:17-18 S '62. (MIRA 17:3)

1. Ukrainskiy nauchno-issledovatel'skiy institut mekhanizatsii
i elektrifikatsii sel'skogo khozyaystva.

KLEYNER, M.K., inzh.; IL'NITSKIY, G.S., inzh.

Heat processes in open-hearth furnaces fired with mixed
gases with carburetted masut. Stal' 24 no.1:27-29 Jh '64.
(MIRA 17:2)

SINKIN, G.P.; IL'NITSKIY, I.I., inzhener, redaktor.

[Vibration damper designed by D.I. Ryshkov] Vibrognitel' konstruktsei
D.I. Ryshkova. Sverdlovsk, Gos. nauchno-tekhn. ind+vo mashinostroit. i
sudostroit. lit-ry [Uralo-Sibirskoe otd-nis] 1953. 27 p. (MLRA 7:7)
(Vibration) (Cutting machines)

IL'NITSKIY, I. I.

Dissertation: "Vibrations Occurring in the Cutting of Metal." Cand Tech Sci, Ural
Polytechnic Inst, Sverdlovsk, 1954. (Referativnyy Zhurnal--Mekhanika, Moscow,
Aug 54)

SO: SUM 393, 28 Feb 1955

IL'NITSKIY, I. I. kandidat tekhnicheskikh nauk.

Causes of natural vibrations in cutting tools. Trudy Ural.
politekh.inst. no.63:37-44 '56. (MIRA 10:2)

(Cutting tools--Vibration)

IL'NITSKIY, I. I.

LOSKUTOV, Vasil'y Vasil'yevich; KUVSHINSKIY, V.V., kandidat tekhnicheskikh nauk, retsentsent; VITENBERG, Yu.R., inzhener, retsentsent; IL'NITSKIY, I.I., kandidat tekhnicheskikh nauk, redaktor; SARAFANNIKOVA, G.A., tekhnicheskij redaktor

[Gear-cutting machines] Zaboronnye stanki. Moskva, Gos. nauchno-tekhn.izd-vo mashinostroit. lit-ry, 1957. 73 p. (Nauchno-populiarnaya biblioteka rabochego stanochnika, no.26) (MLBA 10:6)
(Gear-cutting machines)